

PATHWAYS ADDITIONAL REFERENCES NOT IN MAGAZINE: Summer 2015

Clinical Perspectives

Technologies improve accuracy and reliability of lymphedema measurement

By Margie McNeely

References

1. Armer JM, Stewart BR. A comparison of four diagnostic criteria for lymphedema in a post-breast cancer population. *Lymphatic Research and Biology*. 2005;3(4):208-17. doi:10.1089/lrb.2005.3.208.
2. Ancukiewicz M, Russell TA, Otoole J, Specht M, Singer M, Kelada A et al. Standardized method for quantification of developing lymphedema in patients treated for breast cancer. *Int J Radiat Oncol Biol Phys*. 2011;79(5):1436-43. doi:10.1016/j.ijrobp.2010.01.001.
3. Stanton AW, Northfield JW, Holroyd B, Mortimer PS, Levick JR. Validation of an optoelectronic limb volumeter (Perometer). *Lymphology*. 1997;30(2):77-97.
4. Ward LC, Czerniec S, Kilbreath SL. Quantitative bioimpedance spectroscopy for the assessment of lymphoedema. *Breast Cancer Research and Treatment*. 2009;117(3):541-7. doi:10.1007/s10549-008-0258-010.1007/s10549-008-0090-6;
5. Cornish, B.H., Bunce, I.H., Ward, L.C., Jones, L.C., Thomas, B.J., Bioelectrical impedance for monitoring the efficacy of lymphoedema treatment programmes (1996) *Breast Cancer Research and Treatment*, 38 (2), pp. 169-176. , DOI 10.1007/BF01806671; concordance correlation coefficient to evaluate reproducibility (1989) *Biometrics*, 45 (1), pp. 255-268. , DOI 10.2307/2532051.
6. Ward LC, Czerniec S, Kilbreath SL. Operational equivalence of bioimpedance indices and perometry for the assessment of unilateral arm lymphedema. *Lymphatic Research and Biology*. 2009;7(2):81-5. doi:10.1089/lrb.2008.1027.

Clinical Perspectives

Top 10 Self-Care Tips for Lymphedema

By Lori Radke

References

1. McNeely ML, Peddle CJ, Yurick JL, et al. Conservative and dietary interventions for cancer-related lymphedema: a systematic review and meta-analysis. *Cancer*. 2011;117(6):1136-48.
2. Tidhar D, Hodgson P, Shay C, Towers A. A Lymphedema Self-Management Programme: Report on 30 cases. *Physiotherapy Canada*. 2014; 66(4): 404-412.
3. Vignes S, Porcher R, Arrault M, et al. Long-term management of breast cancer-related lymphedema after intensive decongestive physiotherapy. *Breast Cancer Res Treat*. 2007;101(3):285-90.

4. Paramanandam VS, Roberts D. Weight training is not harmful for women with breast cancer-related lymphoedema: A systematic review. *Journal of Physiotherapy*, 60(3), 136-143.