

## PATHWAYS ADDITIONAL REFERENCES NOT IN MAGAZINE: Winter 2014

### Clinical Perspectives

### Exercise and Lymphedema

By Dorit Tidhar & DeCourcy Squire

#### References

1. Havas E, Parviainen T, Vuorela J, Toivanen J, Nikula T, Vihko V. Lymph flow dynamics in exercising human skeletal muscle as detected by scintigraphy. *Journal of Physiology*. 1997;504.1:233—9.
2. Tartaglione G, Pagan M, Morese R, Cappellini GA, Zappala AR, Sebastiani C, et al. Intradermal lymphoscintigraphy at rest and after exercise: a new technique for the functional assessment of the lymphatic system in patients with lymphoedema. *Nuclear medicine communications*. 2010;31:547–51. doi: 10.1097/MNM.0b013e328338277d.
3. Olszewski WL. Contractility patterns of normal and pathologically changed human lymphatics. *annNYacadsci*. 2002;979:52-63.
4. Casley-Smith JR. Changes in the microcirculation at the superficial and deeper levels in lymphoedema: the effects and results of massage, compression, exercise and benzopyrones on these levels during treatment. *Clin Hemorheol Microcirc*. 2000;23(2-4):335-43. Epub 2001/04/26. PubMed PMID: 11321460.
5. Boris M, Weindorf S, Lasinski S. Persistence of lymphedema reduction after noninvasive complex lymphedema therapy. *Oncology (Williston Park, NY)*. 1997;11(1):99-109; discussion 10, 13-4. Epub 1997/01/01. PubMed PMID: 9115856.
6. Katz-Leurer M BJ. Test–retest reliability of arm volume measurement in women with breast cancer related lymphedema. *J Lymphoedema*. 2012;7:8-13.
7. Godoy Mde F, Pereira MR, Oliani AH, de Godoy JM. Synergic effect of compression therapy and controlled active exercises using a facilitating device in the treatment of arm lymphedema. *Int J Med Sci*. 2012;9(4):280-4. Epub 2012/06/16. doi: 10.7150/ijms.3272. PubMed PMID: 22701334; PubMed Central PMCID: PMC3372933.
8. Moseley AL, Piller NB, Carati CJ. The effect of gentle arm exercise and deep breathing on secondary arm lymphedema. *Lymphology*. 2005;38(3):136-45. Epub 2005/12/16. PubMed PMID: 16353491.
9. McClure MK, McClure RJ, Day R, Brufsky AM. Randomized controlled trial of the Breast Cancer Recovery Program for women with breast cancer-related lymphedema. *The American journal of occupational therapy : official publication of the American Occupational Therapy Association*. 2010;64(1):59-72. Epub 2010/02/06. PubMed PMID: 20131565.
10. Tidhar D, Katz-Leurer M. Aqua lymphatic therapy in women who suffer from breast cancer treatment-related lymphedema: a randomized controlled study. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer*. 2010;18(3):383-92. Epub 2009/06/06. doi: 10.1007/s00520-009-0669-4. PubMed PMID: 19495810.
11. Schmitz KH, Ahmed RL, Troxel AB, Cheville A, Lewis-Grant L, Smith R, et al. Weight lifting for women at risk for breast cancer-related lymphedema: a randomized trial. *JAMA : the journal of the American Medical Association*. 2010;304(24):2699-705. Epub 2010/12/15. doi: 10.1001/jama.2010.1837. PubMed PMID: 21148134.

12. Johansson K, Hayes S, Speck RM, Schmitz KH. Water-based exercise for patients with chronic arm lymphedema: a randomized controlled pilot trial. *American journal of physical medicine & rehabilitation / Association of Academic Physiatrists*. 2013;92(4):312-9. Epub 2013/02/02. doi: 10.1097/PHM.0b013e318278b0e8. PubMed PMID: 23370582.
13. Hayes SC, Johansson K, Stout NL, Prosnitz R, Armer JM, Gabram S, et al. Upper-body morbidity after breast cancer: incidence and evidence for evaluation, prevention, and management within a prospective surveillance model of care. *Cancer*. 2012;118(8 Suppl):2237-49. Epub 2012/04/18. doi: 10.1002/cncr.27467. PubMed PMID: 22488698.
14. Schmitz KH, Ahmed RL, Troxel A, Chevillat A, Smith R, Lewis-Grant L, et al. Weight lifting in women with breast-cancer-related lymphedema. *The New England journal of medicine*. 2009;361(7):664-73. Epub 2009/08/14. doi: 10.1056/NEJMoa0810118. PubMed PMID: 19675330.
15. Kwan ML, Cohn JC, Armer JM, Stewart BR, Cormier JN. Exercise in patients with lymphedema: a systematic review of the contemporary literature. *Journal of cancer survivorship : research and practice*. 2011;5(4):320-36. Epub 2011/10/18. doi: 10.1007/s11764-011-0203-9. PubMed PMID: 22002586.
16. McNeely ML, Campbell KL, Courneya KS, Mackey JR. Effect of acute exercise on upper-limb volume in breast cancer survivors: a pilot study. *Physiotherapy Canada Physiotherapie Canada*. 2009;61(4):244-51. Epub 2010/09/03. doi: 10.3138/physio.61.4.244. PubMed PMID: 20808486; PubMed Central PMCID: PMC2793699.
17. Johansson K, Piller N. Weight bearing exercise and its impact on arm lymphoedema. *Journal of Lymphoedema*. 2007;2(1):15-22.
18. Jonsson C, Johansson K. Pole walking for patients with breast cancer-related arm lymphedema. *Physiotherapy theory and practice*. 2009;25(3):165-73. Epub 2009/04/23. doi: 10.1080/09593980902776621. PubMed PMID: 19384736.
19. Mori T K-LM. Self –measurement of upper extremity volume in women with lymphedema secondary to breast cancer treatment: reliability and validity study: Tel Aviv; 2013.